

Table 1. Northern willowherb control with *granular* herbicides when seeds are applied 0 or 30 days after herbicide application.

Herbicide	Rate (lb/acre)	Seeded 0 days after herbicide application		Seeded 30 days after herbicide application	
		Weed number	SDW (g)	Weed number	SDW (g)
Broadstar	150	10.5	0.0	15.8	0.9
Kansel	100	14.7	7.5	9.0	11.1
OH2	100	12.3	4.2	9.2	0.3
Pendulum	200	12.2	0.4	12.3	0.7
PrePair	150	0.2	0.0	3.5	0.0
Regal O-O	100	9.7	1.3	9.7	1.4
Regal Star	200	8.2	8.2	13.5	0.2
RegalKade	200	9.3	6.7	7.7	2.3
Ronstar	200	1.7	0.0	5.3	0.0
Rout	100	9.2	4.5	13.0	0.4
Snapshot	200	9.5	16.7	10.2	9.0
Non-treated control		4.0	20.4	5.7	4.6

All data collected 60 days after herbicide application.

Table 2. Northern willowherb control with some *granular and sprayed* herbicides when seeds are applied 0 or 30 days after herbicide application.

Herbicide	Rate (lb ai/acre)	Seeded 0 days after herbicides		Seeded 30 days after herbicide application	
		Weed number	SDW (g)	Weed number	SDW (g)
RegalKade	1.5	10.3	14.3	7.7	1.5
Ronstar	4	2.2	3.5	3.0	0.2
Rout	3	9.0	14.2	5.5	0.8
Devrinol + Gallery	1 + 12	11.5	2.6	8.7	1.0
Factor + Gallery	1 + 1.5	8.5	7.6	6.8	0.8
Surflan + Gallery	1 + 4	6.7	0.0	7.8	0.6
Devrinol	12	13.3	1.4	10.0	0.9
Factor	1.5	9.2	11.0	7.2	1.2
Surflan	4	8.0	0.4	8.3	1.4
Gallery	1	14.2	13.4	8.2	2.1
Non-treated control		15.5	15.1	9.3	2.8

All data collected 60 days after herbicide application.